

# Stability of line solitons for KP-II

水町 徹 (九州大学・数理)

2013年7月27日

The KP-II equation

$$\partial_x(\partial_t u + \partial_x^3 u + 3\partial_x(u^2)) + 3\sigma\partial_y^2 u = 0 \quad \text{for } t > 0 \text{ and } (x, y) \in \mathbb{R}^2 \quad (\text{KP})$$

is a generalization to two spatial dimensions of the KdV equation

$$\partial_t u + \partial_x^3 u + 3\partial_x(u^2) = 0 \quad \text{for } t > 0 \text{ and } x \in \mathbb{R}. \quad (1)$$

Obviously, any solutions of (1) are solutions of (KP) which are uniform in  $y$ . The KP equation takes slow variations of the transversal direction into account. If  $\sigma = 1$ , then (KP) is called KP-II and it describes the motion of shallow water waves with weak surface tension.

The KdV equation has a two parameter family of solitary wave solutions

$$\{\varphi_c(x - 2ct - \gamma) \mid c > 0, \gamma \in \mathbb{R}\}, \quad \varphi_c(x) = c \operatorname{sech}^2\left(\sqrt{\frac{c}{2}}x\right).$$

The KP-II equation is expected to explain stability of these solitary wave solutions to the transversal perturbations (see [7]). When  $\varphi_c(x - 2ct)$  is considered as a solution of the KP equations, we call it a *line soliton*. In this talk, I will discuss stability of line soliton solutions for the KP-II equation. The KP-II equation is known to be well-posed in  $L^2(\mathbb{R}^2)$  on the background of line solitons ([11]).

The KP equation (KP) is an integrable system as well as the KdV equation and have conserved quantities such as

$$\int_{\mathbb{R}^2} u(t, x, y)^2 dx dy \quad (\text{momentum}),$$
$$\frac{1}{2} \int (u_x^2(t, x, y) - 3\sigma(\partial_x^{-1}\partial_y u(t, x, y))^2 - 2u^3(t, x, y)) dx dy \quad (\text{Hamiltonian}).$$

For the KP-I equation, that is (KP) with  $\sigma = -1$ , the first two terms of the Hamiltonian have the same sign and it was shown by [6, 14] that there exists a stable ground state for the KP-I equation, whereas the KP-II equation has no traveling wave solutions that belong to  $L^2(\mathbb{R}^2)$  ([5]).

On the other hand, it is known that the line solitons of the KP-I equation are unstable ([16]) and line solitons of the KP-II equation are linearly stable. Because the quadratic part of the Hamiltonian of the KP-II equation, which is the dominant part, is indefinite, it is more natural to explain stability of line solitons by using propagation estimates such as [13] rather than by using variational arguments such as [3].

The main difference between stability analysis of the KdV 1-soliton or the stability of line soliton with the  $y$ -periodic boundary condition is that  $\varphi_c(x - 2ct)$  does not have the finite  $L^2$ -mass because it is not localized in the  $y$ -direction. As a consequence, the linearized operator of the KP-II equation around the line soliton has a family of continuous eigenvalues converging to 0 (see e.g. [1, 2]) in exponentially weighted space, whereas 0 is an isolated eigenvalue of the linearized KdV operator around a solitary wave in exponentially weighted space.

Since we have continuous spectrum converging to 0, the modulations of the speed and the phase shift of line solitons for the KP-II equation cannot be described by ODEs as KdV ([13]) or the KP-II equation posed on  $\mathbb{R}_x \times \mathbb{R}_y / (2\pi\mathbb{Z})$  ([10]).

I find that for the KP-II equation posed on  $\mathbb{R}^2$ , modulation of the speed parameter  $c(t, y)$  and the phase shift  $x(t, y)$  cannot be uniform in  $y$  and their long time behavior is described by the Burgers equation. Note that similar modulation equations have formally derived for a two spatial dimensional Boussinesq model ([12]).

My results are the following ([9]).

**Theorem 1** *Let  $c_0 > 0$  and  $a \in (0, \sqrt{c_0/2})$ . Then there exist positive constants  $\varepsilon_0$  and  $C$  satisfying the following: if  $u(0, x, y) = \varphi_{c_0}(x) + v_0(x, y)$ ,  $v_0 \in H^1(\mathbb{R}^2)$  and  $\varepsilon := \|e^{ax}v_0\|_{L^2(\mathbb{R}^2)} + \|e^{ax}v_0\|_{L_y^1 L_x^2} + \|v_0\|_{L^2(\mathbb{R}^2)} <$*

$\varepsilon_0$ , then there exist  $C^1$ -functions  $c(t, y)$  and  $x(t, y)$  such that for  $t \geq 0$ ,

$$\|u(t, x, y) - \varphi_{c(t, y)}(x - x(t, y))\|_{L^2(\mathbb{R}^2)} \leq C\varepsilon, \quad (2)$$

$$\sup_{y \in \mathbb{R}} (|c(t, y) - c_0| + |x_y(t, y)|) \leq C\varepsilon(1+t)^{-1/2}, \quad (3)$$

$$\|x_t(t, \cdot) - 2c(t, \cdot)\|_{L^2} \leq C\varepsilon(1+t)^{-3/4}, \quad (4)$$

$$\|e^{ax}(u(t, x + x(t, y), y) - \varphi_{c(t, y)}(x))\|_{L^2} \leq C\varepsilon(1+t)^{-3/4}. \quad (5)$$

We find that  $c(t, y)$  and  $\partial_y x(t, y)$  behave like a self-similar solution of the Burgers equation around  $y = \pm\sqrt{8c_0 t}$ .

**Theorem 2** *Let  $c_0 = 2$  and let  $v_0$  and  $\varepsilon$  be the same as in Theorem 1. Then for any  $R > 0$ ,*

$$\left\| \begin{pmatrix} c(t, \cdot) \\ x_y(t, \cdot) \end{pmatrix} - \begin{pmatrix} 2 & 2 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} u_B^+(t, y + 4t) \\ u_B^-(t, y - 4t) \end{pmatrix} \right\|_{L^2(|y \pm 4t| \leq R\sqrt{t})} = o(t^{-1/4})$$

as  $t \rightarrow \infty$ , where  $u_B^\pm$  are self similar solutions of the Burgers equation

$$\partial_t u = 2\partial_y^2 u \pm 4\partial_y(u^2)$$

such that

$$u_B^\pm(t, y) = \frac{\pm m_\pm H_{2t}(y)}{2(1 + m_\pm \int_0^y H_{2t}(y_1) dy_1)}, \quad H_t(y) = (4\pi t)^{-1/2} e^{-y^2/4t},$$

and that  $m_\pm$  are constants satisfying

$$\int_{\mathbb{R}} u_B^\pm(t, y) dy = \frac{1}{4} \int_{\mathbb{R}} c(0, y) dy + O(\varepsilon^2).$$

The KP-II equation (KP) is invariant under a change of variables

$$x \mapsto x + ky - 3k^2 t + \gamma \quad \text{and} \quad y \mapsto y - 6kt \quad \text{for any } k, \gamma \in \mathbb{R}, \quad (6)$$

and has a 3-parameter family of line soliton solutions

$$\mathcal{A} = \{\varphi_c(x + ky - (2c + 3k^2)t + \gamma) \mid c > 0, k, \gamma \in \mathbb{R}\}.$$

Thanks to propagations of the local phase shifts along the crest of line solitons, the set  $\mathcal{A}$  is not stable in  $L^2(\mathbb{R}^2)$ .

**Theorem 3** *Let  $c_0 > 0$ . There exists a positive constant  $C$  such that for any  $\varepsilon > 0$ , there exists a solution of (KP) such that  $\|u(0, x, y) - \varphi_{c_0}(x)\|_{L^2} < \varepsilon$  and*

$$\liminf_{t \rightarrow \infty} t^{-1/4} \inf_{v \in \mathcal{A}} \|u(t, x, y) - v\|_{L^2(\mathbb{R}^2)} \geq C\varepsilon.$$

I remark that similar problems have already been studied for nonlinear heat equations ([15, 8]) and also for kink solutions to a wave equation ([4]). However, we need some correction of the phase shift for the KP-II equation, which does not appear in these former results because the transversal dimension and the power of nonlinearity is lower in our problem.

## 参考文献

- [1] J. C. Alexander, R. L. Pego and R. L. Sachs, *On the transverse instability of solitary waves in the Kadomtsev-Petviashvili equation*, Phys. Lett. A **226** (1997), 187–192.
- [2] S. P. Burtsev, *Damping of soliton oscillations in media with a negative dispersion law*, Sov. Phys. JETP **61** (1985).
- [3] T. Cazenave and P. L. Lions, *Orbital stability of standing waves for some nonlinear Schrödinger equations*, Comm. Math. Phys. **85** (1982), 549–561.
- [4] S. Cuccagna, *On asymptotic stability in 3D of kinks for the  $\phi^4$  model*, Trans. Amer. Math. Soc. **360** (2008), 2581–2614.
- [5] A. de Bouard and Y. Martel, *Non existence of  $L^2$ -compact solutions of the Kadomtsev-Petviashvili II equation*, Math. Ann. **328** (2004) 525–544.
- [6] A. Bouard and J. C. Saut, *Remarks on the stability of generalized KP solitary waves*, Mathematical problems in the theory of water waves, 75–84, Contemp. Math. **200**, Amer. Math. Soc., Providence, RI, 1996.
- [7] B. B. Kadomtsev and V. I. Petviashvili, *On the stability of solitary waves in weakly dispersive media*, Sov. Phys. Dokl. **15** (1970), 539–541.
- [8] T. Kapitula, *Multidimensional stability of planar traveling waves*, Trans. Amer. Math. Soc. **349** (1997), 257–269.
- [9] T. Mizumachi, *Stability of line solitons for the KP-II equation in  $\mathbb{R}^2$* , <http://arxiv.org/abs/1303.3532>

- [10] T. Mizumachi and N. Tzvetkov, *Stability of the line soliton of the KP-II equation under periodic transverse perturbations*, *Mathematische Annalen* **352** (2012), 659–690.
- [11] L. Molinet, J. C. Saut and N. Tzvetkov, *Global well-posedness for the KP-II equation on the background of a non-localized solution*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **28** (2011), 653–676.
- [12] G. Pedersen, *Nonlinear modulations of solitary waves*, *J. Fluid Mech.* **267** (1994), 83–108.
- [13] R. L. Pego and M. I. Weinstein, *Asymptotic stability of solitary waves*, *Comm. Math. Phys.* **164** (1994), 305–349.
- [14] Y. Liu and X. P. Wang, *Nonlinear stability of solitary waves of a generalized Kadomtsev-Petviashvili equation*, *Comm. Math. Phys.* **183** (1997), 253–266.
- [15] J. X. Xin, *Multidimensional stability of traveling waves in a bistable reaction-diffusion equation, I*. *Comm. Partial Differential Equations* **17** (1992), 1889–1899.
- [16] V. Zakharov, *Instability and nonlinear oscillations of solitons*, *JEPT Lett.* **22**(1975), 172-173.